

#27/82  
4-6-01

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE



In re the application of: J. Don Chen

Serial No.: 09/041,994

Filed: March 13, 1998

For: A Transcriptional Coactivator of Steroid/Nuclear  
Receptors and Uses Therefor

Attorney Docket No.: UMG-026 (formerly UMM-026)

Group Art Unit: 1646

Examiner: M. Pak


Assistant Commissioner For Patents  
Washington, D.C. 20231

Certificate of First Class Mailing (37 CFR 1.8(a))

I hereby certify that this correspondence is being deposited with the United States Postal Service as first class mail in an envelope addressed to: Assistant Commissioner for Patents, Washington, D.C. 20231 on the date set forth below.

March 29, 2001  
Date of Signature and of Mail Deposit

By:

  
Debra J. Milasincic  
Reg. No. 46,931  
Attorney for Applicant

DECLARATION UNDER 37 C.F.R. 1.132 BY

DR. J. DON CHEN AND DR. HUI LI

Dear Sir:

We, Dr. J. Don Chen, a citizen of Taiwan, and Dr. Hui Li, a citizen of The People's Republic of China, both residing in the United States, hereby declare as follows:

1. We are co-inventors of the subject matter described and claimed in the above-identified application.

2. We are also co-authors with Paulo J. Gomes of an article entitled "RAC3, a steroid/nuclear receptor-associated coactivator that is related to SRC-1 and TIF2" which was published in *Proc. Natl. Acad. Sci. USA*, vol. 94:8479-8484 (1997), hereinafter, "Li et al."

3. The work described by Li et al. is Applicants' own work. In addition, to the extent that the work described by Li et al. falls within the scope of the claims in the above-identified application, Paulo J. Gomes provided only technical assistance.

We hereby declare that all statements made herein of our own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

3/28/01  
Date

J. Don Chen  
J. Don Chen

3/28/01  
Date

Hui Li  
Hui Li

RECEIVED

APR 03 2001

TECH CENTER 1600/2900

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE



In re the application of: J. Don Chen *et al.*

Serial No.: 09/041,994

Filed: March 13, 1998

For: A Transcriptional Coactivator of Steroid/Nuclear  
Receptors and Uses Therefor

Attorney Docket No.: UMG-026 (formerly UMM-026)

Group Art Unit: 1646

Examiner: M. Pak


Assistant Commissioner for Patents  
Washington, D.C. 20231

Certificate of First Class Mailing (37 CFR 1.8(a))

I hereby certify that this correspondence is being deposited with the United States Postal Service as first class mail in an envelope addressed to: Assistant Commissioner for Patents, Washington, D.C. 20231 on the date set forth below.

March 29, 2001  
Date of Signature and of Mail Deposit

By:

  
Debra J. Milasincic, Esq.  
Reg. No. 46,931  
Attorney for Applicant

DECLARATION UNDER 37 C.F.R. 1.131 BY

DR. J. DON CHEN AND DR. HUI LI

Dear Sir:

We, Dr. J. Don Chen, a citizen of Taiwan, and Dr. Hui Li, a citizen of The People's Republic of China, both residing in the United States, hereby declare as follows:

1. We are co-inventors of the subject matter described and claimed in the above-identified application.

2. Prior to August 5, 1997, the invention described and claimed in the above-referenced application was completed in this country, as evidenced by the following:

- A. Prior to August 5, 1997, we, as co-inventors, isolated a full-length human RAC3 cDNA, determined the nucleic acid sequence of the full-length RAC3 cDNA and derived the human RAC3 amino acid sequence from the RAC3 nucleic acid sequence. The biological activity of RAC3 as a steroid/nuclear receptor-associated coactivator was also determined. We described the full-length RAC3 amino acid sequence and its biological activity in a manuscript entitled "RAC3, a steroid/nuclear receptor-associated coactivator that is related to SRC-1 and TIF2", which was published in *The Proceedings of the National Academy of Sciences, USA* on August 5, 1997. Each of these events is evidenced by a copy of the above-described manuscript (Li *et al.* (1997) *Proc. Natl. Acad. Sci. USA* 94:8479-8484) which is attached hereto as Appendix A, pages A-1 to A-6. We also deposited the full-length RAC3 nucleic acid and amino acid sequences with the GenBank™ sequence database at the National Center for Biotechnology Information (NCBI) prior to August 5, 1997, as evidenced by the attached printout of Genbank™ Accession No. AF010227 (the RAC3 nucleic acid sequence) and AAC51663 (the RAC3 amino acid sequence). Copies of these GenBank™ records are attached hereto as pages A-7 through A-9 and pages A-10 through A-11 of Appendix A.

Thus, the manuscript and GenBank™ Accession Records submitted as pages A-1 through A-11 of Appendix A demonstrate that we had successfully isolated, sequenced, characterized and described human RAC3 nucleic acid and protein molecules, prior to August 5, 1997.

We hereby declare that all statements made herein of our own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the

like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

3/28/01  
Date

J. Don Chen  
J. Don Chen

3/28/01  
Date

Hui Li  
Hui Li

## RAC3, a steroid/nuclear receptor-associated coactivator that is related to SRC-1 and TIF2

HUI LI, PAULO J. GOMES, AND J. DON CHEN\*

Department of Pharmacology and Molecular Toxicology, University of Massachusetts Medical School, 55 Lake Avenue North, Worcester, MA 01655-0126

Communicated by Bert W. O'Malley, Baylor College of Medicine, Houston, TX, June 2, 1997 (received for review April 24, 1997)

**ABSTRACT** Steroids, thyroid hormones, vitamin D<sub>3</sub>, and retinoids are lipophilic small molecules that regulate diverse biological effects such as cell differentiation, development, and homeostasis. The actions of these hormones are mediated by steroid/nuclear receptors which function as ligand-dependent transcriptional regulators. Transcriptional activation by ligand-bound receptors is a complex process requiring dissociation and recruitment of several additional cofactors. We report here the cloning and characterization of receptor-associated coactivator 3 (RAC3), a human transcriptional coactivator for steroid/nuclear receptors. RAC3 interacts with several liganded receptors through a mechanism which requires their respective ligand-dependent activation domains. RAC3 can activate transcription when tethered to a heterologous DNA-binding domain. Overexpression of RAC3 enhances the ligand-dependent transcriptional activation by the receptors in mammalian cells. Sequence analysis reveals that RAC3 is related to steroid receptor coactivator 1 (SRC-1) and transcriptional intermediate factor 2 (TIF2), two of the most potent coactivators for steroid/nuclear receptors. Thus, RAC3 is a member of a growing coactivator network that should be useful as a tool for understanding hormone action and as a target for developing new therapeutic agents that can block hormone-dependent neoplasia.

Steroids, thyroid hormones, vitamin D<sub>3</sub>, and retinoids are fat-soluble small molecules that control cell differentiation, embryonic development, and homeostasis, as well as adult organ physiology (1, 2). The diverse biological effects of these molecules are mediated by a large superfamily of steroid/nuclear hormone receptors which display substantial specificity in regulating gene expression (3–6). These receptors share a common domain structure, including an N-terminal DNA-binding domain (DBD), which binds to specific DNA sequences, and a C-terminal ligand-binding domain (LBD), which binds to the cognate hormones. Retinoic acid receptors (RARs), thyroid hormone receptors (TRs), vitamin D<sub>3</sub> receptor (VDR), peroxisomal proliferator-activated receptors (PPARs), and several other orphan receptors form heterodimeric complexes with retinoid-X receptors (RXRs) (7, 8). Such receptor heterodimers bind to a broad range of response elements, which are composed of two related half-sites and activate target gene expression (9).

Transcriptional activation by steroid/nuclear receptors apparently involves at least two separate processes: derepression and activation (3). Repression is effected in part by association of unliganded receptors with nuclear receptor corepressor (N-CoR) and/or silencing mediator for RAR and TR (SMRT) (10, 11). Ligand binding triggers dissociation of these corepressors and recruitment of coactivators (12). To date, several putative receptor-associated coactivators (RACs) have been

identified, including RIP-140 and RIP-160 (13, 14), ERAP-140 and ERAP-160 (15), TIF1 (16), SRC-1 (17, 18), TRIP1/SUG1 (19, 20), ARA70 (21), TIF2 (22), and CBP/p300 (17, 23, 24). Two of these potential coactivators, SRC-1 (steroid receptor coactivator 1) and TIF2 (transcriptional intermediate factor 2), are related proteins which can enhance transcriptional activation by several steroid/nuclear receptors (18, 22, 24, 25).

Hormone binding is thought to induce conformational changes in the receptor and in turn, activate the C-terminal ligand-dependent activation function (AF-2) of the receptor (2). At the extreme C terminus of the AF-2 domain, about 20 amino acids form an amphipathic helix (26). This helix is referred to as AF-2 activation domain (AF2-AD) (27),  $\tau$ C, or  $\tau$ 4 domain (28, 29). Deletion and several point mutations of this domain abolish the AF-2 function (30–32), and it can act alone as an activation domain when fused to a heterologous DBD (31). Comparison of LBD crystal structures of the unliganded RXR $\alpha$  (26) with liganded RAR $\gamma$  (27) and liganded TR $\alpha$  (33) reveals a striking difference in the relative position of the AF2-AD helix. It is proposed that, upon hormone binding, this helix rotates almost 180° and forms part of the hormone-binding surface. The hydrophobic residues of the helix face toward the ligand-binding cavity, while the charged residues extend into the solvent, possibly mediating protein-protein interactions with cofactors. AF2-AD also has been shown to be required for derepression by promoting dissociation of corepressors (11, 28). These findings suggest a possible mechanistic link among hormone binding, transcriptional repression, and activation by steroid/nuclear receptors through the AF2-AD.

To understand the molecular mechanisms of transcriptional regulation by steroid/nuclear receptors, we screened a human brain cDNA library for RAR-interacting partners by using the yeast two-hybrid system (34). We describe here the cloning and characterization of a novel RAC. RAC3 displays properties of a transcriptional coactivator, including the capacity for ligand-dependent interactions with the receptors and direct transcriptional activation. Interestingly, RAC3 is highly related to SRC-1 and TIF2, suggesting the existence of a family of transcriptional coactivators for steroid/nuclear receptors. The identification of these three related coactivators has provided new insights into the fundamental mechanisms of gene activation by steroids/thyroid hormones, vitamin D<sub>3</sub>, and retinoids.

**Abbreviations:** SRC-1, steroid receptor coactivator 1; TIF2, transcriptional intermediate factor 2; RAC, receptor-associated coactivators; RAR, retinoic acid receptor; RXR, retinoid-X receptor; TR, thyroid hormone receptor; VDR, vitamin D<sub>3</sub> receptor; PPAR, peroxisomal proliferator-activated receptor; PR, progesterone receptor; DBD, DNA-binding domain; LBD, ligand-binding domain; AF, activation function; AF2-AD, AF-2 activation domain; CBP, CREB-binding protein; h, human; m, murine; bHLH, basic helix-loop-helix; PAS, Per-AhR-Sim.

**Data deposition:** The sequence reported in this paper has been deposited in the GenBank database (accession no. AF010227).

\*To whom reprint requests should be addressed. e-mail: J.Chen@ummed.edu.

The publication costs of this article were defrayed in part by page charge payment. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. §1734 solely to indicate this fact.

© 1997 by The National Academy of Sciences 0027-8424/97/948479-6\$2.00/0  
PNAS is available online at <http://www.pnas.org>.

## MATERIALS AND METHODS

**Yeast Two-Hybrid Screen.** The plasmid vector pGBT-hRAR $\alpha$  expressing Gal4 DBD fusion with full-length human RAR $\alpha$  in yeast cells was constructed and used as bait for yeast two-hybrid screening. A human brain cDNA library in pGAD10 vector (CLONTECH) was screened for RAR-interacting proteins in the absence of ligand as previously described (34). After primary selection on synthetic dropout plates lacking tryptophan, leucine, and histidine, but supplemented with 50 mM 3-aminotriazole, we isolated 20 colonies, which were further tested for  $\beta$ -galactosidase expression by using a liquid assay as described (35). The library plasmids from positive clones that expressed both HIS3 and LacZ reporters were rescued and retransformed into yeast cells, together with the original bait and other constructs, for testing the specificity of protein-protein interaction. These analyses led to a single specific clone, RAC3.1, which was selected for further analysis in this study.

**Transient Transfection Assay.** Monkey kidney-derived CV-1 cells and human lung carcinoma A549 cells were grown in phenol red-free Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% charcoal-stripped fetal bovine serum (GIBCO). One day before transfection, cells were seeded in 12-well plates at a density of 50,000 cells per well. A DNA mixture containing 0.5  $\mu$ g of expression vector, 0.5  $\mu$ g of internal control plasmid pCMX- $\beta$ Gal, 1  $\mu$ g of luciferase reporter, and 1.5  $\mu$ g of carrier DNA pGEM was prepared in a final volume of 30  $\mu$ l. The DNA solution was mixed dropwise with 1 vol of 0.5 M CaCl<sub>2</sub> and 2 vol of 2 $\times$  BBS [50 mM N,N-bis(2-hydroxyethyl)-2-aminoethanesulfonic acid (Bes; Calbiochem)/280 mM NaCl/1.5 mM Na<sub>2</sub>HPO<sub>4</sub>, pH 6.95]. DNA precipitates were allowed to form at room temperature for 10 min and applied evenly to the culture cells. Transfection was allowed to continue for 12 hr, and the precipitates were removed by washing transfected cells twice with phosphate-buffered saline (PBS). Transfected cells were re-fed fresh medium containing either vehicle alone or vehicle plus ligands and were harvested 24–36 hr after treatment.

**Luciferase and  $\beta$ -Galactosidase Assay.** Transfected cells in each well were lysed in 120  $\mu$ l of cell lysis solution (35) and processed for luciferase and  $\beta$ -galactosidase assay. Fifty microliters of lysed cells was transferred into 96-well Microtiter plates for luciferase assay and 96-well microtiter plates for  $\beta$ -galactosidase assay as described (35). The luciferase activities were determined with a MLX microtiter plate luminometer (Dynex) using 100  $\mu$ l of assay solution (0.1 M potassium phosphate buffer at pH 7.8, 5 mM ATP, 10 mM MgCl<sub>2</sub>) and 100  $\mu$ l of luciferin solution (0.01 M D-luciferin in 0.1 M potassium phosphate, pH 7.8). The luciferase activities were normalized to the  $\beta$ -galactosidase activity expressed from the cotransfected pCMX- $\beta$ Gal plasmid.

**Isolation and Construction of Full-Length RAC3.** The cDNA insert of RAC3.1 clone was labeled with [<sup>32</sup>P]dATP by DECAprime II DNA labeling kit (Ambion), and used to screen a  $\lambda$ gt11 HeLa cDNA library (CLONTECH). Three overlapping cDNA clones covering the full-length RAC3 coding region were identified and their sequences were determined by dideoxynucleotide sequencing using a T7 sequencing kit (Amersham). The sequence analysis and comparison were carried out with the GCG package. A full-length RAC3 expression vector (pCMX.F.RAC3) was constructed in the pCMX expression vector (9), containing a FLAG and a hemagglutinin epitope linked to the N terminus of RAC3.

## RESULTS

To understand the mechanisms of transcriptional regulation by steroid/nuclear hormone receptors, we screened a human brain cDNA library by the yeast two-hybrid system and identified one specific RAR-interacting protein designated RAC3.

In addition to RAR, RAC3 also interacted with RXR, VDR, PPAR, and TR, but not with COUP-TFI or nonreceptor proteins such as p53 and SNF1 (Fig. 1). In the absence of ligand, RAC3 can interact with RAR and PPAR, and ligand treatment enhanced the RAR interaction by about 75% and PPAR interaction by about 10%. RAC3 also interacted with RXR, VDR, and TR, and these interactions were completely ligand dependent. These results suggest that RAC3 is a specific receptor-interacting protein which preferentially associates with ligand-bound nuclear receptors.

The ligand dependency of the interactions of RAC3 with the receptors suggests that a ligand-induced conformational change can be important for mediating protein-protein interaction with RAC3. Since the AF2-AD domain of the receptors is assumed to undergo a dramatic conformational change upon ligand binding (27), and since this domain is absolutely required for ligand-dependent transcriptional activation (32), we tested three AF2-AD deletion mutants for their abilities to interact with RAC3 in the presence or absence of hormones. Truncation of the C-terminal AF2-AD domain of RAR at residue 403 created the dominant-negative mutant RAR403, which acts as a constitutive repressor in mammalian cells (30, 36), but in yeast cells acted as a strong ligand-dependent transcriptional activator (Fig. 2). The mechanism underlying this difference between yeast and mammalian cells is unclear; however, a similar opposite activation effect by v-erbA in yeast cells was observed (37). Coexpression of RAR403 fusion with RAC3 did not further stimulate the reporter gene activity in either absence or presence of ligand, indicating that the interaction between intact RAR and RAC3 was abolished by deletion of AF2-AD. A similar conclusion was obtained by using mammalian culture cells (data not shown), in which a Gal4 DBD-RAR403 fusion did not activate transcription in response to ligand treatment (35). Truncation of the AF2-AD




FIG. 1. RAC3 interacts with nuclear receptors. The protein-protein interactions between RAC3 and several steroid/nuclear receptors were analyzed by the yeast two-hybrid system. The original yeast two-hybrid clone, RAC3.1 in pGAD10 vector, was retransformed together with yeast expression vectors for Gal4-DBD fusion of selected hormone receptors into Y190 cells. The receptors used include RAR (full-length hRAR $\alpha$ ), RXR (full-length hRXR $\alpha$ ), VDR (LBD of hVDR), PPAR (LBD of mPPAR $\alpha$ ), TR (LBD of hTR $\beta$ ), and COUP (LBD of COUP-TFI) (h-, human; m-, murine). Three independent colonies from each transformation were selected and analyzed for expression of  $\beta$ -galactosidase activity by liquid *o*-nitrophenyl  $\beta$ -D-galactoside (ONPG) assay after treatment with 1  $\mu$ M corresponding ligands (filled bars), or with an equal concentration of vehicle alone (open bars). Under the conditions used in these experiments, ligand treatment did not produce detectable  $\beta$ -galactosidase activity from the Gal4 DBD-receptor fusion alone. atRA, all-trans retinoic acid; VitD3, 1,25-dihydroxyvitamin D<sub>3</sub>; Wy, Wy 14,642; T3, 3,5,3'-triiodothyronine.




FIG. 2. AF2-AD is required for interaction with RAC3. The indicated Gal4-AD fusion (RAC3.1) and Gal4-DBD fusion were cotransformed into yeast Y190 cells, and the  $\beta$ -galactosidase activities from three independent colonies were determined. The yeast cells were treated with 1  $\mu$ M indicated ligands (filled bars) or with an equal concentration of vehicle alone (open bars). RAR403: full-length hRAR $\alpha$  truncated at amino acid 403; RXR443, full-length hRXR $\alpha$  truncated at amino acid 443; VDR364, LBD of hVDR truncated at amino acid 364.

of RXR (RXR443) and VDR (VDR364), unlike that of RAR, did not create ligand-dependent transcriptional activators in yeast cells, while the abilities to interact with RAC3 were totally eliminated (Fig. 2). These results demonstrate that the AF2-ADs of RAR, RXR, and VDR are all absolutely required for interactions with RAC3; thus, RAC3 is an AF-2 dependent cofactor for nuclear receptors.

The above results suggest RAC3 as one of the components of a transcriptionally active complex of liganded receptors. Therefore, we further tested whether RAC3 can directly stimulate transcription when recruited to a specific promoter by linking with a heterologous DBD. Transient transfection of a Gal4 DBD-RAC3 fusion into CV-1 cells resulted in strong stimulation of gene expression from a luciferase reporter containing five copies of Gal4-binding sites, but not from a parental reporter without the Gal4-binding sites (Fig. 3A). The Gal4 DBD alone did not activate the reporter, while a Gal4 DBD fusion of VP16 activation domain strongly stimulated reporter gene expression. Similar experiments in yeast cells also demonstrated a RAC3-dependent transcription stimulation from a *lacZ* reporter in this organism (Fig. 3B). Thus, RAC3 itself contains a transcriptional activation domain functional in both mammalian and yeast cells.

Sequence analysis revealed that the yeast two-hybrid clone RAC3.1 encodes a polypeptide of 804 amino acids (Fig. 4A, in between arrows). Database search revealed a weak similarity of this polypeptide to the central domains of two recently identified steroid receptor coactivators, SRC-1 (17, 18, 38) and TIF2/GRIP1 (22, 39, 40). Full-length RAC3 cDNA was cloned and the nucleotide sequences were determined. The deduced amino acid sequences indicated that RAC3 is a 154.4-kDa protein containing 1417 residues (Fig. 4A). Sequence alignment of full-length RAC3, TIF2 (22), and human SRC-1 (38) revealed a striking homology at the N-terminal region of about 350 amino acids (Fig. 4B). This region contains a potential basic helix-loop-helix (bHLH) domain similar to many transcriptional regulators (41), and two Per-AhR-Sim (PAS) domains found in several nuclear proteins, including Period (Per), aryl hydrocarbon receptor (AhR) and its heterodimeric partner ARNT, the Single Minded protein (Sim), and the hypoxia-inducible factor HIF1 $\alpha$  (42). Sequence comparison among these bHLH-PAS-containing proteins revealed




FIG. 3. RAC3 contains a transcriptional activation domain. The ability of RAC3 to stimulate transcription was tested in both mammalian and yeast cells. (A) Mammalian cells. The Gal4 DBD alone (–) or its fusion with RAC3.1 or VP16 activation domain was expressed in CV-1 cells together with a luciferase reporter containing five copies of Gal4-binding sites (Gal4-tk-Luc) or the parental vector without Gal4-binding sites (tk-Luc). The relative luciferase activities are averages of three independent transfections normalized to  $\beta$ -galactosidase activity. (B) Yeast cells. Three independent colonies expressing Gal4 DBD fusion of RAC3.1 in Y190 cells were analyzed for their  $\beta$ -galactosidase activities. The average  $\beta$ -galactosidase units with standard deviation are shown. The control experiment (–) is  $\beta$ -galactosidase activity of Y190 cells transformed with Gal4 DBD alone.

that RAC3, TIF2, and SRC-1 share a higher degree of similarity than the other proteins (Fig. 4B), suggesting these three receptor-associated coactivators constitute a unique bHLH-PAS family.

Within the RAC3.1 encoding region (amino acid residues 401–1024), we identified six specific motifs sharing a consensus sequence of LXXLL or LLXXL (Fig. 4A). These motifs and their neighboring residues are highly charged and well conserved among RAC3, TIF2, and SRC-1 (Fig. 4C, i to vi). Secondary structure prediction suggests that all these regions have the potential of forming helices, especially motifs iv, v, and vi. Because both transcriptional activation and receptor-interaction activities of RAC3 are located within the RAC3.1 fragment, it is possible that these conserved motifs may play a role in these functions. In addition, a similar motif was found at the C terminus of SRC-1, located within the originally identified receptor-interacting domain (Fig. 4C, vii) (18). At the C-terminal domain of RAC3, a glutamine-rich (Q-rich) domain was identified (Fig. 4A). This Q-rich domain is also conserved among RAC3, TIF2, and SRC-1 and, interestingly, a stretch of 26 consecutive glutamine residues was found only in RAC3. A schematic domain structure comparison among these three related proteins is shown in Fig. 4D.

The ability of RAC3 to physically interact with liganded receptors in an AF-2-dependent manner and to directly stimulate transcription when recruited to a heterologous promoter, together with its sequence similarity to two of the most potent receptor coactivators, are properties which strongly suggest that RAC3 functions as a receptor-associated coactivator. To demonstrate the coactivator function of RAC3, we transiently transfected a mammalian expression vector for full-length RAC3 into human lung carcinoma A549 and/or monkey kidney CV-1 cells. Overexpression of RAC3 enhanced the ligand-dependent transcriptional activation by Gal4 DBD-RAR fusion on a Gal4-dependent luciferase reporter in A549 cells (Fig. 5A), as well as in CV-1 but not in HeLa cells, which express a high level of RAC3 (data not shown). We also tested the effect of overexpression of RAC3 on the transcriptional activation by the human progesterone receptor (hPR $\beta$ ) and found that RAC3 also enhanced transcription by wild-type



observed with full-length human SRC-1 under the test conditions (Fig. 5B). These results demonstrate that RAC3 is




FIG. 5. RAC3 enhances transcriptional activation by steroid/nuclear receptors. (A) The pCMX-F.RAC3 construct expressing full-length RAC3 was transfected into A549 cells together with a construct expressing Gal4-DBD fusion of the LBD of hRAR $\alpha$  (Gal-RAR) and a Gal4-tk-Luc reporter. The relative luciferase activities are averages from three independent experiments after normalization to  $\beta$ -galactosidase activity used as internal controls for transfection efficiency. Transfected cells were treated with vehicle alone (–) or with 100 nM of all-*trans*-retinoic acid for 24 hr after transfection. Overexpression of RAC3 does not have an effect on the luciferase reporter lacking Gal4-binding sites (not shown). (B) RAC3 enhances transcriptional activation by PR on MMTV-LTR promoter. Transfected cells were treated with or without 100 nM progesterone in the presence or absence of coexpressed RAC3 or SRC-1 in CV-1 cells.

capable of enhancing transcriptional activation by liganded receptors in mammalian cells, suggesting that RAC3 is indeed a *bona fide* transcriptional coactivator for steroid/nuclear hormone receptors.

## DISCUSSION

We identified RAC3 as a nuclear receptor-interacting protein on the basis of its ability to interact with RAR in a yeast two-hybrid screen. Several lines of evidence support the position that RAC3 is a transcriptional coactivator for steroid/nuclear hormone receptors. First, we showed that RAC3 interacts with ligand-activated receptors. Second, we demonstrated that the ligand-dependent interaction with RAC3 requires an intact AF2-AD on the receptor. Third, we found that RAC3 itself contains a transcriptional activation domain. Finally, we demonstrated that overexpression of RAC3 enhances ligand-stimulated transcriptional activation by the receptors. In summary, the biochemical properties of RAC3, together with its sequence similarity to SRC-1 and TIF2, identify this protein as a distinct third member of the RAC gene family.

At least two distinct activation functional domains (i.e., AF-1 and AF-2) have been identified in steroid/nuclear hormone receptors (43). The N-terminal AF-1 domain is constitutively active and its activity is not regulated by hormone. In contrast, the activity of the C-terminal AF-2 domain depends on ligand binding. The exact role of the C-terminal AF2-AD helix in recruitment of coactivators is unclear. It is known that the AF2-AD alone is not sufficient for interaction with RAC3 or other coactivators (unpublished data), indicating that other regions of the LBD are also required for recruitment of coactivators. Consistent with this hypothesis, several other activation motifs throughout the LBD of TR have been reported (28). Alternatively, the AF2-AD domain may simply stabilize ligand binding by forming part of the contacting surface for ligands (27, 33), which in turn induces additional

conformational changes that allow association with coactivators.

RAC3 domains responsible for receptor interaction and transcriptional activation remain to be determined. Our data indicated that both domains are located within the central region of RAC3. The receptor-interacting domain of TIF2/GRIP1 has also been located to the central domains (22, 39), while a receptor-interacting domain of SRC-1 was originally found at the C terminus (18). Analysis of the murine SRC-1 has revealed an additional RAR-interacting domain located at the central region (44), where three of the six conserved LXXLL motifs were found. We have confirmed that the N-terminal 782 amino acids of human SRC-1, as well as the C-terminal SRC-1 fragment (amino acids 1245–1440), can interact with several steroid/nuclear hormone receptors in a ligand-dependent manner (unpublished results). However, the C-terminal domain of RAC3 (amino acids 1017–1417), which does not contain any LXXLL motif, cannot interact with liganded receptors (unpublished data). Similar motifs were also found in other potential nuclear receptor coactivators such as RIP-140 (13) and CREB-binding protein (CBP)/p300 (45, 46). These results are consistent with the model that these conserved regions are important for interaction with liganded receptors. Additional domain mapping and mutagenesis experiments should provide further insights into the mechanism of receptor–coactivator interaction. Whether or not some of these conserved motifs are involved in transcriptional activation is also unknown. It is noted that three of these LXXLL motifs are located within the p300-binding domain in mSRC-1 (44); thus, it is possible that these motifs may serve to recruit p300 and/or the basal transcriptional apparatus.

The high degree of sequence conservation within the bHLH-PAS region among RAC3, TIF2, and SRC-1, as compared with other bHLH-PAS proteins, indicates that these three coactivators belong to a distinct family. Intriguingly, the bHLH-PAS domain is dispensable for coactivator functions, including receptor interaction, transcriptional activation, and coactivation (18, 39, 40, 44). The bHLH domain of several transcriptional regulators has been shown to function as a distinct DNA-binding and dimerization motif (42, 47). Whether the bHLH domain in RAC3 and in TIF2 and SRC-1 can mediate protein–protein or protein–DNA interactions is currently unclear. The PAS domains have been shown to be involved in protein–protein interaction in ARNT and its heterodimeric partners (48); however, its role in receptor coactivators is unclear.

SRC-1 has been shown to enhance transcriptional activation by several steroid/nuclear hormone receptors, including PR, growth hormone receptor (GR), estrogen receptor (ER), TR, PPAR, and RXR (18, 25, 49), and it can function synergistically with the common coactivator CBP/p300 (17, 24, 44). It has been proposed that SRC-1 can stabilize a functional interaction between AF-1 and AF-2 of the ER (25). TIF2/GRIP1 has also been shown to enhance transcriptional activities of several steroid/nuclear receptors (22, 40). Our results demonstrated that RAC3 can at least enhance transcriptional activation by RAR and PR. However, it is not clear why we observed only a 2-fold enhancement in both cases by RAC3, while a much higher level of enhancement was observed previously by SRC-1 (18). We reason that this might be due the fact that, in our experiments, an initial 175-fold induction was obtained in PR by hormone treatment in the absence of cotransfected RAC3 or SRC-1, which is in contrast to a 5-fold initial enhancement in an earlier report (18). Therefore, a saturation effect might have limited our ability to detect high level of enhancement. Furthermore, in these experiments we used an SRC-1 cDNA clone that contains the N-terminal bHLH-PAS domain, while in the earlier report a variant of SRC-1 lacking the bHLH-PAS domain was used. By direct comparison between RAC3 and SRC-1, it is clear that RAC3 has a similar

degree of coactivation function like SRC-1 in CV-1 cells under our assay conditions. Further analysis of the effect of RAC3 on transcriptional activation by other steroid/nuclear receptors and a direct comparison with SRC-1 and TIF2 will be necessary to understand how these three related proteins work together to regulate gene activation by hormone receptors.

**Note Added in Proof.** While this paper was in press, Torchia et al. (50) reported a CBP-interacting protein (p/CIP, GenBank accession no. AF000581) which is essential for transcriptional activation by steroid/nuclear receptors as well as by other CBP-dependent transcription factors. RAC3/pCIP shares about 78% identity, including a highly diverged C-terminal 103 amino acids, suggesting that p/CIP is an alternatively spliced form of the murine homologue of RAC3. In the same issue of *Nature*, Heery et al. (51) reported that the first three conserved LXXLL motifs (motifs i, ii, and iii) in SRC-1 are essential and sufficient to mediate protein-protein interactions with liganded receptors.

We thank Drs. E. Bresnick, N. Brown, K. Carlson, W. F. Greenlee, C. L. Peterson, and D. Waud, and other colleagues for stimulating discussion and critical reading of the manuscript. We are very grateful to Dr. K. Umesono for the MMTV-LTR luciferase reporter plasmid and Dr. R. M. Evans for plasmids originated from his laboratory. We also acknowledge Dr. B. O'Malley and M.-J. Tsai for providing the hPR<sub>B</sub> and SRC-1 expression constructs. This work was supported by Institutional Grant IRG203 from the American Cancer Society and a Howard Hughes Pilot Grant for Genetic Research, both provided through the University of Massachusetts Medical School, and an American Society of Hematology Junior Faculty Scholar Award to J.D.C.

1. Evans, R. M. (1988) *Science* **240**, 889–895.
2. Mangelsdorf, D. J., Thummel, C., Beato, M., Herrlich, P., Schütz, G., Umesono, K., Blumberg, B., Kastner, P., Mark, M., Chambon, P. & Evans, R. M. (1995) *Cell* **83**, 835–839.
3. Mangelsdorf, D. J. & Evans, R. M. (1995) *Cell* **83**, 841–850.
4. Beato, M., Herrlich, P. & Schütz, G. (1995) *Cell* **83**, 851–857.
5. Kastner, P., Mark, M. & Chambon, P. (1995) *Cell* **83**, 859–869.
6. Thummel, C. S. (1995) *Cell* **83**, 871–877.
7. Yu, V. C., Delsert, C., Anderson, B., Holloway, J. M., Devary, O. V., Narr, A. M., Kim, S. Y., Boutin, J. M., Glass, C. K. & Rosenfeld, M. G. (1991) *Cell* **67**, 1251–1266.
8. Klierer, S. A., Umesono, K., Mangelsdorf, D. J. & Evans, R. M. (1992) *Nature (London)* **355**, 446–449.
9. Umesono, K., Murakami, K. K., Thompson, C. C. & Evans, R. M. (1991) *Cell* **65**, 1255–1266.
10. Horlein, A. J., Naar, A. M., Heinzel, T., Torchia, J., Gloss, B., Kurokawa, R., Ryan, A., Kamei, Y., Soderstrom, M., Glass, C. K. & Rosenfeld, M. G. (1995) *Nature (London)* **377**, 397–404.
11. Chen, J. D. & Evans, R. M. (1995) *Nature (London)* **377**, 454–457.
12. Fondell, J. D., Ge, H. & Roeder, R. G. (1996) *Proc. Natl. Acad. Sci. USA* **93**, 8329–8333.
13. Cavailles, V., Dauvois, S., L'Horset, F., Lopez, G., Hoare, S., Kushner, P. J. & Parker, M. G. (1995) *EMBO J.* **14**, 3741–3751.
14. Cavailles, V., Dauvois, S., Danielian, P. S. & Parker, M. G. (1994) *Proc. Natl. Acad. Sci. USA* **91**, 10009–10013.
15. Halachmi, S., Marden, E., Martin, G., MacKay, H., Abbondanza, C. & Brown, M. (1994) *Science* **264**, 1455–1458.
16. Le Douarin, B., Zechel, C., Garnier, J. M., Lutz, Y., Tora, L., Pierrat, B., Heery, D., Gronemeyer, H., Chambon, P. & Losson, R. (1995) *EMBO J.* **14**, 2020–2033.
17. Kamei, Y., Xu, L., Heinzel, T., Torchia, J., Kurokawa, R., Gloss, B., Lin, S. C., Heyman, R. A., Rose, D. W., Glass, C. K. & Rosenfeld, M. G. (1996) *Cell* **85**, 403–414.
18. Oñate, S. A., Tsai, S. Y., Tsai, M. J. & O'Malley, B. W. (1995) *Science* **270**, 1354–1357.
19. Lee, J. W., Ryan, F., Swaffield, J. C., Johnston, S. A. & Moore, D. D. (1995) *Nature (London)* **374**, 91–94.
20. vom Baur, E., Zechel, C., Heery, D., Heine, M. J., Garnier, J. M., Vivat, V., Le Douarin, B., Gronemeyer, H., Chambon, P. & Losson, R. (1996) *EMBO J.* **15**, 110–124.
21. Yeh, S. & Chang, C. (1996) *Proc. Natl. Acad. Sci. USA* **93**, 5517–5521.
22. Voegel, J. J., Heine, M. J. S., Zechel, C., Chambon, P. & Gronemeyer, H. (1996) *EMBO J.* **15**, 3667–3675.
23. Chakravarti, D., LaMorte, V. J., Nelson, M. C., Nakajima, T., Schulman, I. G., Juguilon, H., Montminy, M. & Evans, R. M. (1996) *Nature (London)* **383**, 99–103.
24. Smith, C. L., Onate, S. A., Tsai, M. J. & O'Malley, B. W. (1996) *Proc. Natl. Acad. Sci. USA* **93**, 8884–8888.
25. McInerney, E. M., Tsai, M. J., O'Malley, B. W. & Katzenellenbogen, B. S. (1996) *Proc. Natl. Acad. Sci. USA* **93**, 10069–10073.
26. Bourguet, W., Ruff, M., Chambon, P., Gronemeyer, H. & Moras, D. (1995) *Nature (London)* **375**, 377–382.
27. Renaud, J. P., Rochel, N., Ruff, M., Vivat, V., Chambon, P., Gronemeyer, H. & Moras, D. (1995) *Nature (London)* **378**, 681–689.
28. Banihmad, A., Leng, X., Burris, T. P., Tsai, S. Y., Tsai, M. J. & O'Malley, B. W. (1995) *Mol. Cell. Biol.* **15**, 76–86.
29. Hollenberg, S. M. & Evans, R. M. (1988) *Cell* **55**, 899–906.
30. Damm, K., Heyman, R., Umesono, K. & Evans, R. M. (1993) *Proc. Natl. Acad. Sci. USA* **90**, 2989–2993.
31. Barettoni, D., Vivanco Ruiz, M. d. M. & Stunnenberg, H. G. (1994) *EMBO J.* **13**, 3039–3049.
32. Durand, B., Saunders, M., Gaudon, C., Roy, B., Losson, R. & Chambon, P. (1994) *EMBO J.* **13**, 5370–5382.
33. Wagner, R. L., Aprelletti, J. W., West, B. L., Baxter, J. D. & Fletterick, R. J. (1995) *Nature (London)* **378**, 690–697.
34. Durfee, T., Becherer, K., Chen, P. L., Yeh, S. H., Yang, Y., Kilburn, A. E., Lee, W. H. & Elledge, S. J. (1993) *Genes Dev.* **7**, 555–569.
35. Chen, J. D., Umesono, K. & Evans, R. M. (1996) *Proc. Natl. Acad. Sci. USA* **93**, 7567–7571.
36. Tsai, S. & Collins, S. J. (1993) *Proc. Natl. Acad. Sci. USA* **90**, 7153–7157.
37. Privalsky, M. L., Sharif, M. & Yamamoto, K. R. (1990) *Cell* **63**, 1277–1286.
38. Takeshita, A., Yen, P. M., Misiti, S., Cardona, G. R., Liu, Y. & Chin, W. W. (1996) *Endocrinology* **137**, 3594–3597.
39. Hong, H., Kohli, K., Trivedi, A., Johnson, D. L. & Stallcup, M. R. (1996) *Proc. Natl. Acad. Sci. USA* **93**, 4948–4952.
40. Hong, H., Kohli, K., Garabedian, M. J. & Stallcup, M. R. (1997) *Mol. Cell. Biol.* **17**, 2735–2744.
41. Murre, C., McCaw, P. S. & Baltimore, D. (1989) *Cell* **56**, 777–783.
42. Swanson, H. I. & Bradfield, C. A. (1993) *Pharmacogenetics* **3**, 213–230.
43. Tasset, D., Tora, L., Fromental, C., Scheer, E. & Chambon, P. (1990) *Cell* **62**, 1177–1187.
44. Yao, T. P., Ku, G., Zhou, N., Scully, R. & Livingston, D. M. (1996) *Proc. Natl. Acad. Sci. USA* **93**, 10626–10631.
45. Arany, Z., Sella, W., Livingston, D. M. & Eckner, R. (1994) *Cell* **77**, 799–800.
46. Chrivia, J. C., Kwok, R. P. S., Lamb, N., Hagiwara, M., Montminy, M. R. & Goodman, R. H. (1993) *Nature (London)* **365**, 855–859.
47. Murre, C., McCaw, P. S., Vaessin, H., Caudy, M., Jan, L. Y., Jan, Y. N., Cabrera, C. V., Buskin, J. N., Hauschka, S. D., Lassar, A. B., Weintraub, H. & Baltimore, D. (1989) *Cell* **58**, 537–544.
48. Lindebro, M. C., Poellinger, L. & Whitelaw, M. L. (1995) *EMBO J.* **14**, 3528–3539.
49. DiRenzo, J., Söderström, M., Kurokawa, R., Ogliastro, M.-H., Ricote, M., Ingrey, S., Hörlein, A., Rosenfeld, M. G. & Glass, C. K. (1997) *Mol. Cell. Biol.* **17**, 2166–2176.
50. Torchia, J., Rose, D. W., Inostroza, J., Kamei, Y., Westin, S., Glass, C. K. & Rosenfeld, M. G. (1997) *Nature (London)* **387**, 677–684.
51. Heery, D. M., Kalkhoven, E., Hoare, S. & Parker, M. G. (1997) *Nature (London)* **387**, 733–736.

LOCUS AF010227 4495 bp mRNA PRI 13-AUG-1997  
DEFINITION Homo sapiens receptor-associated coactivator 3 (RAC3) mRNA,  
complete cds.  
ACCESSION AF010227  
VERSION AF010227.1 GI:2318005  
KEYWORDS .  
SOURCE human.  
ORGANISM Homo sapiens  
Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;  
Mammalia; Eutheria; Primates; Catarrhini; Hominidae; Homo.  
REFERENCE 1 (bases 1 to 4495)  
AUTHORS Li, H., Gomes, P.J. and Chen, J.D.  
TITLE RAC3, a steroid/nuclear receptor-associated coactivator that is  
related to SRC-1 and TIF2  
JOURNAL Proc. Natl. Acad. Sci. U.S.A. 94 (16), 8479-8484 (1997)  
MEDLINE 97385128  
REFERENCE 2 (bases 1 to 4495)  
AUTHORS Chen, J.D., Li, H. and Gomes, P.J.  
TITLE Direct Submission  
JOURNAL Submitted (25-JUN-1997) Pharmacology and Molecular Toxicology,  
University of Massachusetts Medical School, 55 Lake Avenue North,  
Worcester, MA 01655-0126, USA  
FEATURES  
source Location/Qualifiers  
1..4495  
/organism="Homo sapiens"  
/db\_xref="taxon:9606"  
/cell\_line="HeLa"  
gene 1..4495  
/gene="RAC3"  
CDS 86..4339  
/gene="RAC3"  
/note="transcriptional coactivator with intrinsic  
>histone acetyltransferase activity; member of the  
steroid/nuclear receptor-associated coactivator family  
which includes steroid receptor coactivator 1 (SRC-1),  
transcriptional intermediate factor 2 (TIF2), and receptor  
associated coactivator 3; similar to the mouse  
p300/CBP/co-integrator protein (p/CIP) and the activator  
of retinoid receptors (ACTR)"  
/codon\_start=1  
/product="receptor-associated coactivator 3"  
/protein\_id="AAC51663.1"  
/db\_xref="GI:2318006"  
/translation="MSG LGENLDPLASDSRKRKLP CDT PGQGLTCSGEKRRREQESKY  
IEELAE LISANLSDIDNFNVKPKCAILKETVRQIRQIKEQGKTI SNDDDVQKADVSS  
TGQGVIDKDSLGLPLLQALDGLFVFNREANIVFVSENV TQYLQYKQEDLVNTSVYNI  
LHEEDRKDFLKNLPKSTVNGVSWTNETQRQKSHTFNCRLMKTPHDILEDINASPEMR  
QRYETMQCFALSQPRAMMEEGEDLQSCMICVARRITTGERTFPSNPESFITRHDLSGK  
VVNIDTNSLRSSMRPGFEDIIRRCIQRFSLNDGQSWSQKRHYQEAYLNNGHAETPVYR  
FSLADGTIVTAQTKSKLFRNPVTNDRHGFVSTHFLQREQNGYRPNPNPVGQGI RPPMA  
GCNSSVGGMSMSPNQGLQMPSSRAYGLADPSTTGQMSGARYGGSSNIASLT PGPGMQS  
PSSYQNNYGLNMSSPPHGS PGLAPNQNNIMISPRNRGSPKIASHQFSPVAGVHSPMA  
SSGNTGNHSFSSSSLSALQAISEGVGTSLLSTLSSPGPKLDNSPNMNITQPSKVSNDQ  
SKSPLGIFYCDQNPVESSMCQNSRDHLSDKESKESSVEGAENQRGP LKSGHKKLLQL  
LTCSSDDRGHSSLTNSPLDSSCKESSVSVTSPSGVSSSTSGGVSSTSNMHGSL LQEKH  
RILHKLLQNGNSPAEVAKITAEATGKDTSSITSCGDGNVVKQEQLSPKKKENNALLRY  
LLDRDDPSDALSKELQPQVEGVDNKMSQCTSSTIPSSSQEKDPKIKTETSEEGSGDLD  
NLDAILGDLTSSDFYNNSISSNGSHLGTKQVVFQGTNSLGLKSSQVSQIRPPYNRAV  
SLDSPVSVGSSPPVKNISAFPM LPKQ PMLGGNPRMMDSQENYGS SMGGPNRNVTVTQT  
PSSGDWGLPNSKAGRMEPMNSNSMGRPGGDYNTSLPRPALGGS IPTLPLRSNSIPGAR  
PVLQQQQMLQMRPGEIPMGMGANPYGQAAASNQLGSWPDGMLSMEQVSHGTQNRPLL  
RNSLDDLVGPPSNLEGQSDERALLDQLHTLLSNTDATGLEEIDRALGIPELVNQGGAL  
EPKQDAFQGGQEAAMMDQKAGLYGQTYPAQGPMPQGGFHLQGS PPSFNSMMNQMNQGG  
NFP LQGMHPRANIMRPTNTPKQLRMQLQORLQGGQFLNQSRQALELKMENPTAGGAA  
VMRPMMPQGGFLNAQMVAQRSRELLSHHFRQQRVAMMMQQQQQQQQQQQQQQQQQQQQQ

QQQQQQQTQAFSPPPNVTASPSMDGLLAGPTMPQAPPQQFPYQPNYGMGQQPDPAFGR  
 VSSPPNAMMSSRMGPSQNPMMQHPQAASIYQSSEMKGWPSGNLARNSSFSQQQFAHQG  
 NPAVYSMVHMGSSGHMGQNMNMPMPMSGMPMGPDQKYC"

BASE COUNT  
 ORIGIN

1354 a 1048 c 1043 g 1050 t

1	gctggatggt	ggactcagag	accaataaaa	ataaactgct	tgaacatcct	ttgactgggt
61	agccagttgc	tgatgtatat	tcaagatgag	tggattagga	gaaaacttgg	atccactggc
121	cagtgattca	cgaaaacgca	aattgccatg	tgatactcca	ggacaaggct	ttacctgcag
181	tggtgaaaaa	cggagacggg	agcaggaaa	taaatatatt	gaagaattgg	ctgagctgat
241	atctgccaat	cttagtgata	ttgacaat	caatgtcaaa	ccagataaat	gtgcgatttt
301	aaaggaaaaca	gtaagacaga	tacgtcaaat	aaaagagcaa	ggaaaaacta	tttccaatga
361	tgatgatggt	caaaaaacccg	atgtatcttc	tacagggcag	ggagttattg	ataaagactc
421	cttaggaccg	cttttacttc	aggcatggga	tggtttccta	tttgtgggtga	atcgagaggc
481	aaacatttga	tttgtatcag	aaaatgtcac	acaatacctg	caatataagc	aagaggacct
541	ggttaacaca	agtgtttaca	atatcttaca	tgaagaagac	agaaaggatt	ttcttaagaa
601	tttaccaaaa	tctacagtta	atggagtttc	ctggacaaat	gagacccaaa	gacaaaaaag
661	ccatacat	aattgccgta	tggtgatgaa	aacaccacat	gatattctgg	aagacataaa
721	cgccagtcct	gaaatgcgcc	agagatatga	aacaatgcag	tgctttgccc	tgtctcagcc
781	acgagctatg	atggaggaag	gggaagattt	gcaatcttgt	atgatctgtg	tggcacgccg
841	cattactaca	ggagaaaaga	catttccatc	aaaccctgag	agctttatta	ccagacatga
901	tctttcagga	aaggttgta	atatagatac	aaattcactg	agatcctcca	tgaggcctgg
961	ctttgaagat	ataatccgaa	ggtgtattca	gagatttttt	agtctaaatg	atgggcagtc
1021	atgggtcccag	aaacgtcact	atcaagaagc	ttatcttaat	ggccatgcag	aaacccagct
1081	atatcgattc	tcgttggtcg	atggaactat	agtactgca	cagacaaaaa	gcaaactctt
1141	ccgaaatcct	gtaacaaatg	atcgacatgg	ctttgtctca	accacttcc	ttcagagaga
1201	acagaatgga	tatagaccaa	acccaaatcc	tggtggacaa	gggattagac	cacctatggc
1261	tggatgcaac	agttcggtag	gcggcatgag	tatgtcgcca	aaccaaggct	tacagatgcc
1321	gagcagcagg	gcctatggct	tggcagaccc	tagcaccaca	gggcagatga	gtggagctag
1381	gtatgggggt	tccagtaaca	tagcttcatt	gaccctggg	ccaggcatgc	aatcaccatc
1441	ttcctaccag	aacaacaact	atgggtctca	catgagtgc	ccccacatg	ggagtcctgg
1501	tcttgcccca	aaccagcaga	atatcatgat	ttctcctcgt	aatcgtggga	gtccaaagat
1561	agcctcacat	cagttttctc	ctgttgacag	tgtgcactct	cccatggcat	cttctggcaa
1621	tactgggaac	cacagctttt	ccagcagctc	tctcagtgcc	ctgcaagcca	tcagtgaagg
1681	tgtggggact	tcccttttat	ctactctgtc	atcaccaggc	cccaaattgg	ataactctcc
1741	caatatgaat	attacccaac	caagtaaagt	aagcaatcag	gattccaaga	gtcctctggg
1801	cttttattgc	gacaaaaatc	cagtggagag	ttcaatgtgt	cagtcaaata	gcagagatca
1861	cctcagtgc	aaagaaaagta	aggagagcag	tggtgagggg	gcagagaatc	aaaggggtcc
1921	tttgaaaagc	aaaggtcata	aaaaattact	gcagttactt	acctgttctt	ctgatgaccg
1981	gggtcattcc	tccttgacca	actccccctc	agattcaagt	tgtaaagaat	cttctgttag
2041	tgtcaccagc	ccctctggag	tctcctctc	tacatctgga	ggagtatcct	ctacatccaa
2101	tatgcattgg	tactgtttac	aagagaagca	ccggattttg	cacaagtgc	tgcagaatgg
2161	gaattcacca	gctgaggtag	ccaagattac	tgcagaagcc	actgggaaag	acaccagcag
2221	tataacttct	tgtggggacg	gaaatgttgt	caagcaggag	cagctaagtc	ctaagaagaa
2281	ggagaataat	gcacttctta	gatacctgct	ggacagggat	gatcctagt	atgcactctc
2341	taaagaacta	cagccccaag	tggaaaggat	ggacaataaa	atgagtcagt	gcaccagctc
2401	caccattcct	agctcaagtc	aagagaagga	ccctaaaatt	aagacagaga	caagtgaagc
2461	gggtatctgga	gacttggaata	atctatagtc	tattcttggg	gatctgacta	gttctgactt
2521	ttacaataat	tccatatcct	caaatggtag	tcatctgggg	actaagcaac	aggtgtttca
2581	aggaactaat	tctctgggtt	tgaaaagtgc	acagtctgtg	cagtctattc	gtcctccata
2641	taaccgagca	gtgtctctgg	atagccctgt	ttctgttggc	tcaagtcctc	cagtaaaaaa
2701	tatcagtgtc	ttccccatgt	taccaaagca	acccatgttg	ggtgggaatc	caagaatgat
2761	ggatagtcag	gaaaattatg	gctcaagtat	gggtgggcca	aaccgaaatg	tgactgtgac
2821	tcagactcct	tcctcaggag	actggggctt	accaaactca	aaggccggca	gaatggaacc
2881	tatgaattca	aactccatgg	gaagaccagg	aggagattat	aatacttctt	taccagacc
2941	tgcactgggt	ggctctattc	ccacattgcc	tcttcggtct	aatagcatac	caggtgcgag
3001	accagttatg	caacagcagc	agcagatgct	tcaaatgagg	cctggtgaaa	tccccatggg
3061	aatgggggct	aatccctatg	gccaagcagc	agcatctaac	caactgggtt	cctggcccga
3121	tggcatgttg	tccatggaac	aagtttctca	tggcactcaa	aataggcctc	ttcttaggaa
3181	ttccctggat	gatcttgttg	ggccaccctc	caacctggaa	ggccagagt	acgaaagagc
3241	attattggac	cagctgcaca	ctcttctcag	caacacagat	gcgacaggcc	tggaaagaat
3301	tgacagagct	ttgggcattc	ctgaacttgt	caatcaggga	caggcattag	agcccaaaca
3361	ggatgctttc	caaggccaag	aagcagcagt	aatgatggat	cagaaggcag	gattatatgg
3421	acagacatac	ccagcacagg	ggcctccaat	gcaaggaggc	tttcatcttc	agggacaatc
3481	accatctttt	aactctatga	tgaatcagat	gaaccagcaa	ggcaattttc	ctctccaagg
3541	aatgcaccca	cgagccaaca	tcatgagacc	ccggacaaac	acccccaagc	aacttagaat

3601 gcagcttcag cagaggctgc agggccagca gtttttgaat cagagccgac aggcacttga  
3661 attgaaaatg gaaaacccta ctgctggtgg tgctgcggtg atgaggccta tgatgcagcc  
3721 ccagcagggg tttcttaatg ctcaaatggt cgcccaacgc agcagagagc tgctaagtca  
3781 tcacttccga caacagaggg tggctatgat gatgcagcag cagcagcagc agcaacagca  
3841 gcagcagcag cagcagcagc aacagcaaca gcaacagcag caacagcagc aaaccagggc  
3901 cttcagccca cctcctaata tgactgcttc ccccagcatg gatgggcttt tggcaggacc  
3961 cacaatgcc caagctcctc cgcaacagtt tccatatcaa ccaaattatg gaatgggaca  
4021 acaaccagat ccagcctttg gtcgagtgtc tagtcctccc aatgcaatga tgcgtcaag  
4081 aatgggtccc tcccagaatc ccatgatgca acaccgcag gctgcatcca tctatcagtc  
4141 ctcaaaaatg aagggtctggc catcaggaaa tttggccagg aacagctcct tttcccagca  
4201 gcagtttgcc caccagggga atcctgcagt gtatagtatg gtgcacatga atggcagcag  
4261 tggtcacatg ggacagatga acatgaaccc catgcccatt tctggcatgc ctatgggtcc  
4321 tgatcagaaa tactgctgac atctctgcac caggacctct taaggaaacc actgtacaaa  
4381 tgacactgca ctaggattat tgggaaggaa tcattgttcc aggcattccat cttggaagaa  
4441 aggaccagct ttgagctcca tcaagggtat ttttaagtgt gtcatttgag cagga

A-10

LOCUS AAC51663 1417 aa PRI 12-AUG-1997  
 DEFINITION receptor-associated coactivator 3 [Homo sapiens].  
 ACCESSION AAC51663  
 PID g2318006  
 VERSION AAC51663.1 GI:2318006  
 DBSOURCE locus AF010227 accession AF010227.1  
 KEYWORDS .  
 SOURCE human.  
 ORGANISM Homo sapiens  
 Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;  
 Mammalia; Eutheria; Primates; Catarrhini; Hominidae; Homo.  
 REFERENCE 1 (residues 1 to 1417)  
 AUTHORS Li, H., Gomes, P.J. and Chen, J.D.  
 TITLE RAC3, a steroid/nuclear receptor-associated coactivator that is  
 related to SRC-1 and TIF2  
 JOURNAL Proc. Natl. Acad. Sci. U.S.A. 94 (16), 8479-8484 (1997)  
 MEDLINE 97385128  
 REFERENCE 2 (residues 1 to 1417)  
 AUTHORS Chen, J.D., Li, H. and Gomes, P.J.  
 TITLE Direct Submission  
 JOURNAL Submitted (25-JUN-1997) Pharmacology and Molecular Toxicology,  
 University of Massachusetts Medical School, 55 Lake Avenue North,  
 Worcester, MA 01655-0126, USA  
 COMMENT Method: conceptual translation.  
 FEATURES Location/Qualifiers  
 source 1..1417  
 /organism="Homo sapiens"  
 /db\_xref="taxon:9606"  
 /cell\_line="HeLa"  
 Protein 1..1417  
 /product="receptor-associated coactivator 3"  
 CDS 1..1417  
 /gene="RAC3"  
 /coded\_by="AF010227.1:86..4339"  
 /note="transcriptional coactivator with intrinsic~>histone  
 acetyltransferase activity; member of the steroid/nuclear  
 receptor-associated coactivator family which includes  
 steroid receptor coactivator 1 (SRC-1), transcriptional  
 intermediate factor 2 (TIF2), and receptor associated  
 coactivator 3; similar to the mouse p300/CBP/co-integrator  
 protein (p/CIP) and the activator of retinoid receptors  
 (ACTR) "  
 ORIGIN  
 1 msglgenldp lasdsrkrkl pcdtpggglt csgekrreq eskyieelae lisanlsdid  
 61 nfnvkdpc k ilketvrqir qikeggktis ndddvqkad sstgggvidk dslgplllqa  
 121 ldgflfvvnr eanivfvsen vtqylqykqe dlvntsvyni lheedrkdfi knlpkstvng  
 181 vswtnetqrq kshtfncrml mktphdiled inaspemrqr yetmqcfals qprammeege  
 241 dlqscmicva rrittgertf psnpesfitr hdlsgkvvni dtnslrssmr pgfediirrc  
 301 iqrrffslndg qswsqkrhyq eaylnghaet pvyrfsladg tivtaqtksk lfrnptvndr  
 361 hgfvtsthflq reqngyrpnp npvgggirpp magcnssvgg msmspnqglq mpssraygla  
 421 dpsttgqmsg aryggssnia sltpgpgmq sssyqnnnyg lnmsspphgs pglapnqqni  
 481 misprnrgsp kiashqfspv agvhspmass gntgnhsfss ssalsalqais egvgtsllst  
 541 lsspgpkldn spnmnitqps kvsngqdsks lgyfcdqnpv essmcqsnsr dhlsdkeske  
 601 ssvegaenqr gpleskghkk llqlitcssd drghssltns pldssckess vsvtspsgvs  
 661 sstsggvsst snmhgsllqe khrlhklilq ngnsapaevak itaeatgkdt ssitscgdgn  
 721 vvkqeqlspk kennallry lldrddpsda lskelqpqve gvdnkmsqct sstipsssqe  
 781 kdpkiktets eegsgldlnd dailgdltss dfynnsissn gshlgtkqqv fqgtnsllglk  
 841 ssqsvqsirp pynravsls psvsgssppv knisafpmlp kqpmllgnpr mmdsqenygs  
 901 smggpnrvnt vtqtpssgdw glpnsakgrm epmnsnsmgr pggdyntslp rpalggsipt  
 961 plrsnsipg arpvlqqqqq mlqmnpgeip mgmganpygq aaasnqlgsw pdgmllsmeqv  
 1021 shgtqnrpll rnslddlvgp psnlegqsde ralldqlhtl lsntdatgle eidralgipe  
 1081 lvnqqgalep kqdafggqea avmmdqkagl ygqtypagqp pmqggfhlqg qspfsnsmmn  
 1141 qmnqqgnfpl qgmhpranim rptntpkql rmqlqrlqg qqflnqsrqa lelkmnpta  
 1201 ggaavmrpmm qpqqgflnaq mvagrsrell shhfrqqrva mmmqqqqqqq qqqqqqqqqq  
 1261 qqqqqqqqqt qafspppnvt aspmdglla gptmpqappq qfpyqpnygm gqqpdpafgr

A-11

1321 vssppnamms srmgpsqnpm mqhpqaasiy qssemkgwps gnlarnssfs qqqfahqgnp  
1381 avysmvhmng ssghmgqnm npmpmsgmpm gpdqkyc

//